arXiv:1605.03682v1 [math.GR] 12 May 2016

Effective quasimorphisms on free chains

Jing Tao*

Abstract

We find homogeneous counting quasimorphisms that are effective at seeing chains in a free group **F**. As corollary, we derive that if a group G has an index–d free subgroup, then every element $g \in G$ either has stable commutator length at least $1/8d$ or some power of g is conjugate to its inverse. We also show that for a finitely-generated free group **F**, there is a countable basis for the real vector space of homogeneous quasimorphisms on **F**.

1 Introduction

Counting quasimorphisms on free groups were first constructed by Brooks in [Bro81]. A useful variation of the Brooks' construction was introduced by Epstein and Fujiwara in [EF97]. Their variation counts the number of disjoint copies of a reduced word in another word (see Section 3). This induces a large family of homogeneous counting quasimorphisms on a free group with defect at most 4 [Cal09]. Further, like the Brooks' quasimorphisms, this family *sees* every element of **F**; that is, for any nontrivial element $g \in \mathbf{F}$, there is a word w whose associated homogeneous counting quasimorphism ϕ_w has $\phi_w(g) \geq 1$. Via Bavard's duality, this yields a lower bound of $1/8$ for the stable commutator length (scl) of any nontrivial element in a free group.

In this paper, we are interested in quasimorphisms that see chains of a free group **F**. Let $C_1(\mathbf{F})$ be the vector space of real 1–chains of **F** and let $\overline{C}_1(\mathbf{F})$ be the quotient of $C_1(\mathbf{F})$ by the subspace spanned by elements of the form $g^n - ng$, and $g - hgh^{-1}$, where $g, h \in G$ and $n \in \mathbf{Z}$. Using a lexicographical order on **F**, we define the notion of *effective words* in **F**. Each effective word is indivisible (not power of another word) and is a unique representative of its own conjugacy class. We show that the set of counting quasimorphisms associated to effective words enjoy nice properties. In particular, for every nontrivial chain of the form $\sum_i g_i \in \overline{C}_1(\mathbf{F})$, there is an effective word w whose associated homogeneous counting quasimorphism ϕ_w sees $\sum_i g_i$ (see Proposition 3.3). Using Generalized Bavard's Duality [Cal09] (see Theorem 2.1), we obtain

Theorem 1.1. *Let* **F** *be any free group and let* $g_1, \ldots, g_n \in \mathbf{F}$ *be a finite sequence such that* $\sum_i g_i$ *is a* 1*–boundary. Then, either* $\sum_i g_i$ *is the zero chain in* $\overline{C}_1(\mathbf{F})$*, in which case* $\mathrm{scl}\left(\sum_i g_i\right) = 0$*, or* $\mathrm{scl}\left(\sum_i g_i\right) \geq 1/8$.

Theorem 1.1 has applications to the stable commutator lengths in virtually free groups. Recall a group is *virtually free* if it contains a free subgroup of finite index. Using the finite-index formula of [Cal09] (see Theorem 2.2), we derive the following corollary of Theorem 1.1.

*Partially supported by NSF DMS-1311834.

Corollary 1.2. *Let G be a virtually free group with a free subgroup of index d. Then* $\mathrm{scl}(g) = 0$ *if and only if there exists $n \in \mathbf{Z}$ such that g^n is conjugate to g^{-n}. Otherwise,* $\mathrm{scl}(g) \geq 1/8d$.

Since virtually free groups are δ–hyperbolic, it already follows from [EF97] (see also [CF10]) that there is always a positive lower bound for the scl of every element which does not have a power that is conjugate to its own inverse. Their lower bound depends only on δ but it is not explicit. The advantage of Corollary 1.2 is the explicit lower bound relating the index of the free subgroup.

Finally, we construct a basis for the real vector space of homogeneous quasimorphisms on $\mathbf{F}$ using effective words. More precisely,

Theorem 1.3. *For every free group $\mathbf{F}$, there is a collection of effective words $B \subset \mathbf{F}$ such that the associated set $\{\phi_w\}_{w\in B}$ of homogeneous counting quasimorphisms is linearly independent. Further, for every homogeneous quasimorphism ϕ on $\mathbf{F}$, there are real numbers $\{r_w\}_{w\in B} \subset \mathbf{R}$ such that $\phi = \sum_{w\in B} r_w \phi_w$.*

We remark that the set B has the property that distinct elements represent distinct conjugacy classes, and for every indivisible element $g \in B$, either the conjugacy class of g or the conjugacy class of its inverse is represented in B. As corollary of Theorem 1.3, we obtain that for any countable free group, there is a countable basis for the real vector space of homogeneous quasimorphisms. However, it is essential that infinitely many of the corresponding coefficients r_w are allowed to be nonzero, as there are homogeneous quasimorphisms that are not a finite linear combination of counting quasimorphism. See the end of the paper for an example or Examples 2.63 and 2.64 of [Cal09].

Finally, we note that there is a sharp lower bound of $1/2$ for the scl of nontrivial elements in free groups. This was first shown in [DH91]; (see also [LW14] for another proof). The proof for this lower bound does not use quasimorphisms. Thus, it is natural to ask if the lower bound of Theorem 1.1 can be improved using a different argument.

Acknowledgments

This paper grew out of a joint project with Talia Fernós and Max Forester. I am grateful to Max for many stimulating conversations related to this paper. I also thank Mladen Bestvina for helpful comments and for suggesting the example of the last section. Finally, I thank the referee for thorough and helpful comments.

2 Preliminaries

We refer to [Cal09] and references within for background on quasimorphisms and stable commutator lengths.

Let G be any group. A real-valued function $\Phi\colon G \to \mathbf{R}$ is a *quasimorphism* on G if there exists $D \geq 0$ such that

$$|\Phi(gh) - \Phi(g) - \Phi(h)| \leq D, \quad \forall g, h \in G.$$

The infimum over all such constants D is the *defect* D_Φ of Φ. It is immediate that Φ is a homomorphisms if and only if D_Φ is 0. A quasimorphism is *homogeneous* if it is a homomorphism on cyclic subgroups of G. Homogeneous quasimorphisms are (homogeneous) *class functions*, i.e. they are constant on conjugacy classes of G. Given a quasimorphism Φ, the *homogenization* of Φ is defined by

$$\phi(g) = \lim_{n\to\infty} \frac{\Phi(g^n)}{n}.$$

This is a homogeneous quasimorphism with defect $D_\phi \le 2D_\Phi$ [Cal09, Lemma 2.58].

Let $C_n(G)$ be the real vector space freely generated by n–tuples of elements in G with boundary map $\partial\colon C_n(G) \to C_{n-1}(G)$ defined on each n–tuple by

$$\partial(g_1,\dots,g_n) = \sum_{i=1}^{n} (-1)^{i-1}(g_1,\dots,g_{i-1},g_{i+1},\dots,g_n).$$

A quasimorphism on G extends linearly to $C_1(G)$ and a homogeneous quasimorphism descends to the quotient $\overline{C}_1(G)$ of $C_1(G)$ by the subspace spanned by elements of the form $g^n - ng$, and $g - hgh^{-1}$, where $g,h \in G$ and $n \in \mathbf{Z}$. We denote by $B_1(G) = \partial C_2(G)$ the subspace of real 1–boundaries of G and let $\overline{B}_1(G)$ be the image of $B_1(G)$ in $\overline{C}_1(G)$. For simplicity, we will take the Generalized Bavard's Duality as the definition of the *stable commutator length* (scl) of an element in $\overline{B}_1(G)$. The definition of the scl of a chain (Definition 2.71) and a proof of the duality theorem (Theorem 2.79) can be found in [Cal09].

Theorem 2.1 (Generalized Bavard's Duality)**.** *Let G be any group. Then for any $\sum_i^n r_i g_i \in \overline{B}_1(G)$,*

$$\mathrm{scl}\left(\sum_i^n r_i g_i\right) = \sup_\phi \frac{\sum_i^n r_i \phi(g_i)}{2D_\phi},$$

where ϕ ranges over all homogeneous quasimorphisms on G.

In the sum above, if $n = 1$ and $r_1 = 1$, then we obtain the usual Bavard's duality for $\mathrm{scl}(g)$ of an element $g \in [G,G]$.

We now describe the finite-index formula stated in [Cal09, Proposition 2.80]. Suppose H is a finite-index subgroup of G. Let X be a space with fundamental group G, and let $\widehat{X}$ be the covering space associated to the subgroup H. Given a loop γ representing the conjugacy class of $g \in G$, the preimage in $\widehat{X}$ can be expressed as a finite union of loops $\widehat{\gamma}_1,\dots,\widehat{\gamma}_k$ each covering γ with some finite degree. Suppose $\widehat{\gamma}_i$ represents the conjugacy class of $h_i \in H$. Then the following formula holds.

Theorem 2.2 (Finite index formula)**.**

$$\mathrm{scl}_G(g) \;=\; \frac{1}{[G:H]}\,\mathrm{scl}_H\left(\sum_{i=1}^{k} h_i\right).$$

3 Effective words

Let $\mathbf{F}$ be a free group, possibly with infinite rank. Fix a well-ordered, symmetric, free generating set for $\mathbf{F}$ and equip $\mathbf{F}$ with the lexicographical order. Let $|w|$ be the word length of $w \in \mathbf{F}$. By *conjugacy length* $|w|_C$ of w we will mean the word length of a cyclically reduced representative of the

conjugacy class of w. For each cyclically reduced word $w \in \mathbf{F}$, let $[w]$ be the set of conjugates of w obtained by cyclically permuting the letters of w. Note that $[w]$ is a finite set. We will call a word $w \in \mathbf{F}$ *effective* if w is indivisible and cyclically reduced, and it has minimal lexicographical order among all elements in $[w]$. For any indivisible element $g \in \mathbf{F}$, we will say w is an *effective representative* of g if w is conjugate to g and w is effective.

Let w be a reduced word. For any $g \in \mathbf{F}$, let $C_w(g)$ be the maximum number of disjoint copies of w in the reduced representative of g. Define

$$\Phi_w(g) = C_w(g) - C_{w^{-1}}(g).$$

By [Cal09, Proposition 2.30], Φ_w is a quasimorphism with defect at most 2, thus its homogenization ϕ_w has defect at most 4.

Lemma 3.1. *If $w \in \mathbf{F}$ is effective, then $\phi_w(g) = 0$ for any $g \in \mathbf{F}$ with $|g|_C < |w|$.*

Proof. In this proof, we will denote by $w < w'$ to mean w is smaller than w' lexicographically. We may assume g is cyclically reduced, so $|g| = |g|_C$. Suppose there exists $n > 0$ such that $C_w(g^n) = 1$. Since $|g| < |w|$, we must have $n \geq 2$. By cyclically permuting the letters of g, we may assume $w = (uv)^{n-1}u$, where $g = uv$. By assumption, g is cyclically reduced so $|uv| = |vu|$. We can't have $uv = vu$, since this would imply u and v are both powers of the same word, which contradicts that w is indivisible. Hence, either $uv < vu$ or $vu < uv$. If $uv < vu$, then $uuv < uvu$. But uvu is a prefix of w and uuv is a prefix of $uwu^{-1} = u(uv)^{n-1}$, so $uwu^{-1} < w$, contradicting the minimality of the order of w in $[w]$. On the other hand, if $vu < uv$, then $u^{-1}wu = v(uv)^{n-2}uu < w$, since uv is a prefix of w but vu is a prefix of $u^{-1}wu$. Again, this contradicts minimality of w. We conclude $C_w(g^n) = 0$ for all $n > 0$. Since this is true for all g with $|g| < |w|$, we also have $C_{w^{-1}}(g^n) = C_w((g^{-1})^n) = 0$, for all $n > 0$. This shows $\Phi_w(g^n) = 0$ for all $n \in \mathbf{Z}$, whence the result. □

The following lemma is immediate.

Lemma 3.2. *Suppose $w \in \mathbf{F}$ is cyclically reduced and $g \in \mathbf{F}$ has $|w| = |g|_C$. Then*

- *If g is conjugate to w, then $\phi_w(g) = 1$.*
- *If g is conjugate to w^{-1}, then $\phi_w(g) = -1$.*
- *In all other cases, $\phi_w(g) = 0$.*

The following proposition, together with Theorem 2.1 and the fact that $D_{\phi_w} \leq 4$ for any reduced word $w \in \mathbf{F}$, yields Theorem 1.1.

Proposition 3.3. *For any finite sequence $g_1, \ldots, g_n \in \mathbf{F}$, if $\sum_i g_i$ is not the zero chain in $\overline{C}_1(\mathbf{F})$, then there is an effective word $w \in \mathbf{F}$ with $\sum_i \phi_w(g_i) \geq 1$.*

Proof. Suppose the chain $c = \sum_i g_i$ is nonzero in $\overline{C}_1(\mathbf{F})$. For each i, set $g_i = w_i^{m_i}$, where w_i is indivisible and $m_i \geq 1$. For $j \neq k$, if $w_j = t w_k t^{-1}$, then $g_j + g_k \equiv g_j + t g_k t^{-1} \equiv (m_j + m_k) w_j$ in $\overline{C}_1(\mathbf{F})$. Similarly, if $w_j^{-1} = t w_k t^{-1}$ and $m_j > m_k$, then $g_j + g_k \equiv (m_j - m_k) w_j$. Therefore, by applying these equivalence relations whenever possible, we may assume $c \equiv \sum_j m_j w_j$, where w_j is not conjugate to w_k or w_k^{-1} whenever $j \neq k$ and that each $m_j > 0$. Finally, by replacing g_j by a conjugate if necessary, we will assume each w_j is effective.

By reordering the indices if necessary, we may assume $|w_1| \geq |w_2| \geq \cdots \geq |w_n|$. Let $1 \leq k \leq n$ be the largest index such that $|w_1| = |w_2| = \cdots = |w_k|$. For each i, set $\phi_i = \phi_{w_i}$. By definition, if $k < n$, then $|w_k| > |w_{k+1}| \geq \cdots \geq |w_n|$. Therefore, by Lemma 3.1, for all $1 \leq i \leq k$ and all $j > k$, $\phi_i(w_j) = 0$. Also, by Lemma 3.2, for all $1 \leq i, j \leq k$, $\phi_i(w_j) = 0$ if $i \neq j$, and $\phi_i(w_i) = 1$. Hence, for each $1 \leq i \leq k$,

$$\phi_i(c) = \sum_j m_j \phi_i(w_j) = m_i \geq 1.$$

Thus choosing $w = w_i$ for any $1 \leq i \leq k$ gives the statement of the proposition. □

We remark that the proof also shows that for any nonzero chain $c = \sum_i r_i g_i \in \overline{C}_1(\mathbf{F})$, there is an effective word $w \in \mathbf{F}$ with $\phi_w(c) > 0$.

Proof of Corollary 1.2. Let G be a virtually free group with a free subgroup H of index d. By applying the finite index formula (Theorem 2.2) and Theorem 1.1 to H, we obtain that if $\mathrm{scl}_G(g) > 0$, then $\mathrm{scl}(g) \geq 1/8d$. It remains to show that $\mathrm{scl}_G(g) = 0$ if and only if some power of g is conjugate to its inverse. Though this follows from known results ([EF97] and [CF10]), we will give an independent proof that depends only on our results on chains.

If some power of g is conjugate to its inverse then $\mathrm{scl}_G(g) = 0$ follows from homogeneity. Conversely, suppose $\mathrm{scl}_G(g) = 0$. Let $N \lhd G$ be a normal subgroup of finite index contained in H. The group N is is free, being a subgroup of a free group, so fix a well-ordered, symmetric, free generating set for N. Applying the finite-index formula to N, we obtain that

$$\mathrm{scl}_G(g) \;=\; \frac{1}{[G:N]} \mathrm{scl}_N\left(\sum_{i=1}^{k} h_i\right),$$

where each $h_i \in N$ is conjugate (in G) to some power of g. Since N is normal, each h_i is conjugate to the same power of g. Therefore, it suffices to find a pair h_i and h_j such that h_i is conjugate (in N) to h_j^{-1}. By Theorem 1.1, $\mathrm{scl}_G(g) = 0$ if and only $\sum_i h_i$ is trivial in $\overline{C}_1(N)$. For each i, there exists an effective word $w_i \in N$ and $m_i > 0$ such that $h_i \equiv m_i w_i$ in $\overline{C}_1(N)$. By reordering the indices if necessary, we may assume $|w_1| \geq |w_i|$ for all i. If no w_j is conjugate (in N) to w_1^{-1}, then by Lemma 3.1 and Lemma 3.2, $\sum_i \phi_{w_1}(h_i) = m_1 > 0$, contradicting $\mathrm{scl}_G(g) = 0$. Therefore, w_1 must be conjugate to the inverse of w_j for some j. Since N is normal, the conjugacy lengths $|h_i|_C$ are all the same. Since each w_i is cyclically reduced, $|h_i|_C = m_i|w_i|$. Therefore, $m_1|w_1| = m_j|w_j|$ which implies $m_1 = m_j$. This shows that h_1 is conjugate to the inverse of h_j as desired. □

Basis

Let $\widehat{B}$ be the set of effective words of $\mathbf{F}$. Since an effective representative is unique for each conjugacy class, distinct elements of $\widehat{B}$ represent distinct conjugacy classes. We decompose $\widehat{B} = B \sqcup \overline{B}$, such that B contains the effective representative of g if and only if $\overline{B}$ contains the effective representative of g^{-1}. Note that B forms a basis for $\overline{C}_1(\mathbf{F})$. We impose a well ordering $<$ on B (not the lexicographical order) as follows. For $w, w' \in B$, if $|w| < |w'|$, then $w < w'$; if $|w| = |w'|$, then set $w < w'$ if w is smaller lexicographically. Let $\mathscr{B} = \{\phi_w : w \in B\}$. We show

Lemma 3.4. *Only finitely many elements of $\mathscr{B}$ do not vanish on any nontrivial element $g \in \mathbf{F}$.*

Proof. For any nontrivial element $g \in \mathbf{F}$, there is a unique element in B that has a power conjugate to g or g^{-1}. Thus, by homogeneity, it is enough to show the statement for $g \in B$. For any reduced word w with $|w| < |g|$, if $\phi_w(g) \neq 0$, then w must be a subword of g^2. Since there are only finitely many subwords in g^2, $\phi_w(g) = 0$ for all but finitely many $w \in B$ with $|w| < |g|$. Now by Lemma 3.1 and Lemma 3.2, with the exception of $w = g$, any $w \in B$ with $|w| \geq |g|$ has $\phi_w(g) = 0$. The conclusion follows. □

Let $\mathscr{C}$ be the real vector space of all homogeneous class functions on $\mathbf{F}$. Theorem 1.3 is immediate from the following proposition.

Proposition 3.5. *Any $\phi \in \mathscr{C}$ can be represented by $\phi = \sum_{w \in B} r_w \phi_w$, for some $\{r_w\} \subset \mathbf{R}$. Further, $\mathscr{B}$ is linearly independent.*

Proof. We start with the first statement. For any $w \in B$, denote by $S_w = \{w' \in B : \ w' < w\}$. We now use transfinite induction to define r_w for all $w \in B$. For the base case, let $w_0 \in B$ be the minimal element and set $r_{w_0} = \phi(w_0)$. Now for any $w \in B$, suppose $r_{w'}$ is defined for all $w' \in S_w$. Set

$$r_w = \phi(w) - \sum_{w' \in S_w} r_{w'} \phi_{w'}(w).$$

By Lemma 3.4, the sum in the definition of r_w is well defined since there are only finitely many nonzero terms. This defines r_w for all $w \in B$. To see $\phi = \sum_w r_w \phi_w$, by homogeneity it is enough to show that the two sides agree on every element in B. By Lemma 3.1 and Lemma 3.2, for any $w, g \in B$ with $w \geq g$, $\phi_w(g) = 1$ if and only if $w = g$. Therefore,

$$\sum_w r_w \phi_w(g) = \sum_{w \in S_g} r_w \phi_w(g) + r_g = \phi(g).$$

We now show linearly independence of $\mathscr{B}$. Suppose there exists $\phi_w \in \mathscr{B}$ which is in the span of $\mathscr{B} \setminus \{\phi_w\}$. We first show that $r_{w'} = 0$ for all $w' \neq w$ with $|w'| \leq |w|$. Fix $k \leq |w|$ suppose by induction $r_{w'} = 0$ for all $|w'| < k$. Write

$$\phi_w = \sum_{\substack{|w'| \geq k \\ w' \neq w}} r_{w'} \phi_{w'}.$$

Now let $g \in B \setminus \{w\}$ with $|g| = k \leq |w|$. By Lemma 3.1 and Lemma 3.2, for any $|w'| \geq |g|$, $\phi_{w'}(g) = 1$ if and only if $w' = g$ and is 0 for any other case. Therefore, evaluating on g, we obtain

$$0 = \phi_w(g) = r_g \phi_g(g) + \sum_{\substack{|w'| \geq k \\ w' \neq w, g}} r_{w'} \phi_{w'}(g) = r_g.$$

This shows $r_{w'} = 0$ for all $|w'| \leq |w|$. We now have $\phi_w = \sum_{|w'| > |w|} r_{w'} \phi_{w'}$. Now plug in w into the two sides, we obtain

$$1 = \phi_w(w) = \sum_{|w'| > |w|} r_{w'} \phi_{w'}(w) = 0.$$

This finishes the proof. □

Corollary 3.6. *Let $\mathbf{F}$ be a countable free group. Then the set of class functions with rational coefficients is dense in $\mathscr{C}$ in the operator norm; that is, for any $\phi \in \mathscr{C}$ and $\epsilon > 0$, there are rational numbers $\{q_w\}_{w \in B} \subset \mathbf{Q}$ such that*

$$\frac{\left|\phi(g) - \sum_w q_w \phi_w(g)\right|}{|g|} < \epsilon, \quad \forall g \in \mathbf{F} \setminus \{\mathrm{id}\}.$$

Consequentially, the set of homogeneous quasimorphisms with rational coefficients is dense in the vector space of homogeneous quasimorphisms (in the operator norm or in the defect norm).

Proof. Since $\mathbf{F}$ is countable, so is B. Let $\phi = \sum_w r_w \phi_w \in \mathscr{C}$ be arbitrary and let $\epsilon > 0$ be given. We can choose rational numbers $\{q_w\}_{w\in B} \subset \mathbf{Q}$ such that $\sum_w |r_w - q_w| < \epsilon$. For any $g \in \mathbf{F}$, $|\phi_w(g)| \leq |g|$. Therefore

$$\left|\sum_w \phi_w(g) - \sum_w q_w \phi_w(g)\right| \leq \sum_w \left|r_w - q_w\right| \left|g\right| < \epsilon \left|g\right|.$$

For the second statement, recall that the defect of each ϕ_w is at most 4. Thus, if $\phi = \sum_w r_w \phi_w$ is a quasimorphism, then we also have that for $\psi = \phi - \sum_w q_w \phi_w$ and for all $g, h \in \mathbf{F}$,

$$\left|\psi(gh) - \psi(g) - \psi(h)\right| \leq \sum_w |r_w - q_w| \left|\phi_w(gh) - \phi_w(g) - \phi_w(h)\right| \leq 4\epsilon.$$

This shows $\sum_w q_w \phi_w$ is bounded in the defect norm from ϕ, hence it is a quasimorphism. □

We observe that since each ϕ_w has defect at most 4, if $\sum_w |r_w| < \infty$, then $\phi = \sum_w r_w \phi_w$ is a quasimorphism with defect at most $4\sum_w |r_w| < \infty$. However, a quasimorphism may have non-summable coefficients as illustrated by the following example.

Let $\mathbf{F}_2 = \langle a, b\rangle$ be the free group of rank 2. Order its generators by $a < a^{-1} < b < b^{-1}$. Choose the set B to contain the words a and $a^n b$ for all $n \geq 0$.

For any $g \in \mathbf{F}_2$, represent g in reduced form by $g = a^{p_1} b^{q_1} \cdots a^{p_k} b^{q_k}$, where all the exponents are non-zero integers except for possibly p_1 or q_k. Define $\Phi(g)$ to be the number of positive even exponents minus the number of negative even exponents in the expression for g. The map Φ is a quasimorphism by [Rol09]. Let ϕ be its homogenization and set $\phi = \sum_{w\in B} r_w \phi_w$. We will show by induction that for all $n \geq 2$, $r_{a^n b} = (-1)^n$. This immediately implies that $\sum_{w\in B} |r_w|$ is not summable.

First note that

$$\phi(a) = \phi(b) = \phi(ab) = 0 \quad \Longrightarrow \quad r_a = r_b = r_{ab} = 0.$$

Also, for any $w \in B$, $\phi_w(a^n b) \neq 0$ if and only if $w = a$ or $w = a^m b$ for $0 \leq m \leq n$. In the latter case, $\phi_{a^m b}(a^n b) = 1$. Thus, for all $n \geq 2$, we have

$$r_{a^n b} = \phi(a^n b) - \sum_{w' < a^n b} r_{w'} \phi_{w'}(a^n b) = \phi(a^n b) - \sum_{m=2}^{n-1} r_{a^m b}$$

This establishes the base case $r_{a^2 b} = \phi(a^2 b) = 1$. We now suppose $n > 2$ and for all $2 \leq m < n$, $r_{a^m b} = (-1)^m$. Therefore,

$$r_{a^n b} = \phi(a^n b) - \sum_{m=2}^{n-1} (-1)^m.$$

If n is odd, then $\phi(a^n b) = 0$, and $\sum_{m=2}^{n-1} (-1)^m = 1$. If n is even, then $\phi(a^n b) = 1$, and $\sum_{m=2}^{n-1} (-1)^m = 0$. This yields $r_{a^n b} = (-1)^n$ for all $n \geq 2$ as desired.

References

[Bro81] Robert Brooks. Some remarks on bounded cohomology. In *Riemann surfaces and related topics: Proceedings of the 1978 Stony Brook Conference (State Univ. New York, Stony Brook, N.Y., 1978)*, volume 97 of *Ann. of Math. Stud.*, pages 53–63. Princeton Univ. Press, Princeton, N.J., 1981.

[Cal09] Danny Calegari. *scl*, volume 20 of *MSJ Memoirs*. Mathematical Society of Japan, Tokyo, 2009.

[CF10] Danny Calegari and Koji Fujiwara. Stable commutator length in word-hyperbolic groups. *Groups Geom. Dyn.*, 4(1):59–90, 2010.

[DH91] Andrew J. Duncan and James Howie. The genus problem for one-relator products of locally indicable groups. *Math. Z.*, 208(2):225–237, 1991.

[EF97] David B. A. Epstein and Koji Fujiwara. The second bounded cohomology of word-hyperbolic groups. *Topology*, 36(6):1275–1289, 1997.

[LW14] Larsen Louder and Henry Wilton. Stackings and the W-cycle conjecture. Preprint, 2014. Available at `arXiv:1410.2540 [math.GR]`.

[Rol09] Pascal Rolli. Quasi-morphisms on free groups. Preprint, 2009. Available at `arXiv:0911.4234 [math.GR]`.

Mathematics Department, University of Oklahoma, Norman, OK 73019, USA

`jing@math.ou.edu`